

General Guidelines for Teachers and Staff**

- Suicide is the second leading cause of death for Children between the ages of 10-14, and third leading cause of death for Adolescents between 15-24 years of age in the United States. ***
- In recent years, more young people have died from suicide than from cancer, heart disease, HIV/AIDS, congenital birth defects, and diabetes combined. *
- For every young person who dies by suicide, between 100-200 attempt suicide. *
- Males are four times as likely to die by suicide as females, although females attempt suicide three times as often as males. *

SUICIDE IS PREVENTABLE

Here is what you can do:

- **Talk** to your student about suicide. Asking direct questions will not put ideas into their heads, as many fear. Asking for help is the single greatest skill that will protect your student. Help your student identify and connect to caring adults to talk to when they need guidance and support.
- **Know** the risk factors and warning signs of suicide. Participate and frequently review the suicide prevention training information from your school district.
- **Remain calm.** Becoming too excited or distressed will communicate that you are not able to talk about suicide.
- **Listen** without judging. (practice safe messaging) Allow for the discussion of experiences, thoughts, and feelings. Be prepared for expression of intense feelings. Try to understand the reasons for considering suicide without taking a position about whether such behavior is justified.
- **Supervise** constantly. Do not leave the individual alone until a caregiver (often a parent) or school crisis team member has been contacted and agrees to provide appropriate supervision.
- **Ask** if there is a plan for suicide. If it does not put you or the student in danger, attempt to remove the suicide means.
- **Respond** immediately. Escort the student to a member of your school's crisis team. If you are unsure of who is on your school crisis team, find the

LA County RESOURCE
877.7.CRISIS or 877.727.4747
Suicide Prevention Center

<https://preventsuicide.lacoe.edu>

NATIONAL RESOURCE
800.273.TALK (8255)
National Suicide
Prevention Lifeline

Copyright 2010 by Los Angeles Unified School District

Permission is granted for reproduction and distribution for educational purposes with appropriate acknowledgement of authorship to the Los Angeles Unified School District (LAUSD).

Adapted from: LAUSD and updated 2024 by Los Angeles County Office of Education- Center for Distance and Online Learning (CDOL).

principal, assistant principal, school social worker, psychologist, counselor, or school nurse.

- **Join** the crisis team. You know your student the best. Provide essential background information that will help with assessing the student's risk for suicide. When a teacher says, "this behavior is not like this student", this is critical information indicating a sudden change in behavior.

Youth Suicide Risk Factors

While the path that leads to suicidal behavior is long and complex and there is no "profile" that predicts suicidal behavior with certainty, there are certain risk factors associated with increased suicide risk. In isolation, these factors are not signs of suicidal thinking. However, when present they signal the need to be vigilant for the warning signs of suicide. In addition, they are also appropriate targets for suicide prevention programs. Specifically, these risk factors include the following:

- History of depression, mental illness or substance/alcohol abuse disorders
- Presence of a firearm or rope
- Isolation or lack of social support
- Situational crisis
- Family history of suicide or suicide in community
- Hopelessness
- Impulsivity
- Incarceration
- History of self-injury

Assembly Bills 2246 and 1767 also identify the following high risk groups

- Youth bereaved by suicide
- Youth with Disabilities, mental illness or substance use disorders
- Youth experiencing homelessness or in out of home settings, such as foster care
- Lesbian, gay, bisexual, transgender or questioning youth

LA County RESOURCE
877.7.CRISIS or 877.727.4747
Suicide Prevention Center

<https://preventsuicide.lacoe.edu>

NATIONAL RESOURCE
800.273.TALK (8255)
National Suicide
Prevention Lifeline

Copyright 2010 by Los Angeles Unified School District

Permission is granted for reproduction and distribution for educational purposes with appropriate acknowledgement of authorship to the Los Angeles Unified School District (LAUSD).

Adapted from: LAUSD and updated 2024 by Los Angeles County Office of Education- Center for Distance and Online Learning (CDOL).

Suicide Warning Signs

Warning signs are observable behaviors that may signal the presence of suicidal thinking. They might be considered cries for help or invitations to intervene. These warning signs signal the need to inquire directly about whether the individual has thoughts of suicide. If such thinking is acknowledged, then suicide interventions will be required. Warning signs include the following:

- **Suicide threats.** It has been estimated that up to 80% of all suicide victims have given some clues regarding their intentions. Both direct ("I want to kill myself.") and indirect ("I wish I could fall asleep and never wake up.") threats need to be taken seriously. Consider social media posts that include direct or indirect threats.
- **Suicide notes and plans.** The presence of a suicide note is an especially significant sign of danger. The greater the planning revealed by the youth, the greater the risk of suicidal behavior.
- **Prior suicidal behavior.** Prior behavior is a powerful predictor of future behavior. Thus, anyone with a history of suicidal behavior should be carefully observed for future suicidal behavior.
- **Making final arrangements.** Making funeral arrangements, writing a will, and/or giving away prized possessions may be warning signs of impending suicidal behavior.
- **Preoccupation with death.** Excessive talking, drawing, reading, writing, or posting on social media about death may suggest suicidal thinking.
- **Changes in behavior, appearance, thoughts, and/or feelings.** Depression (especially when combined with hopelessness), sudden happiness (especially when preceded by significant depression), a move toward social isolation, giving away personal possessions, and reduced interest in previously important activities are among the changes considered to be suicide warning signs.

*M. Heron, D. L. Hoyert, S. L. Murphy, J. Xu, K. D. Kochanek, & B. Tejada-Vera. (2009, April). Deaths: Final Data for 2006. National Vital Statistics Reports, 57(14).

LA County RESOURCE
877.7.CRISIS or 877.727.4747
Suicide Prevention Center

<https://preventsuicide.lacoe.edu>

NATIONAL RESOURCE
800.273.TALK (8255)
National Suicide
Prevention Lifeline

Copyright 2010 by Los Angeles Unified School District

Permission is granted for reproduction and distribution for educational purposes with appropriate acknowledgement of authorship to the Los Angeles Unified School District (LAUSD).

Adapted from: LAUSD and updated 2024 by Los Angeles County Office of Education- Center for Distance and Online Learning (CDOL).

**Lieberman, R., Poland, S. & Cassel, R. (2008). Suicide intervention. In Thomas, A. & Grimes, J., Best practices in school psychology V. Bethesda, MD: National Association of School Psychologists.

***WISQARS Leading Causes of death Visualization Tool, CDC, 2022 results

LA County RESOURCE
877.7.CRISIS or 877.727.4747
Suicide Prevention Center

<https://preventsuicide.lacoe.edu>

NATIONAL RESOURCE
800.273.TALK (8255)
National Suicide
Prevention Lifeline

Copyright 2010 by Los Angeles Unified School District
Permission is granted for reproduction and distribution for educational purposes with appropriate
acknowledgement of authorship to the Los Angeles Unified School District (LAUSD).
Adapted from: LAUSD and updated 2024 by Los Angeles County Office of Education- Center for Distance and Online
Learning (CDOL).